

(19)

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 400 243 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
24.03.2004 Bulletin 2004/13

(51) Int Cl.7: **A61K 31/415, A61K 31/33,
A61P 29/00**

(21) Application number: **03255860.3**

(22) Date of filing: **18.09.2003**

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IT LI LU MC NL PT RO SE SI SK TR**
Designated Extension States:
AL LT LV MK

(30) Priority: **19.09.2002 JP 2002272662**
14.03.2003 JP 2003070298
24.07.2003 JP 2003278699

(71) Applicant: **TANABE SEIYAKU CO., LTD.**
Chuo-ku, Osaka-shi, Osaka 541-8505 (JP)


(72) Inventors:
• **Kono, Rikako, Towa City Coop**
Saitama-shi Saitama 330-0854 (JP)

• **Kohnomi, Shuntarou**
Toda-shi Saitama 335-0015 (JP)
• **Aihara, Hajime**
Saitama-shi Saitama 337-0051 (JP)
• **Hosaka, Toshihiro**
Tokyo 110-0015 (JP)
• **Kashiwagi, Toshihiko**
Toda-shi Saitama 335-0015 (JP)

(74) Representative: **Cresswell, Thomas Anthony**
J.A. KEMP & CO.
14 South Square
Gray's Inn
London WC1R 5JJ (GB)

(54) **Calcium-activated K channel activator**

(57) There is disclosed the use of a compound of formula (I):



or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the prophylaxis or treatment of a disorder or disease responsive to opening of the large conductance calcium-activated K channel.

EP 1 400 243 A1

Description

BACKGROUND OF THE INVENTION

1. FIELD OF THE INVENTION

[0001] This invention relates to an excellent large conductance calcium-activated K channel opener, which is useful for treatment of disorders or diseases such as pollakiuria, urinary incontinence, cerebral infarction, subarachnoid hemorrhage, and the like.

2. BACKGROUND ART

[0002] Potassium is the most abundant intracellular cation, and is very important in maintaining physiological homeostasis. Potassium channels are present in almost all vertebrate cells, and the potassium influx through these channels is indispensable for maintaining hyperpolarized resting membrane potential.

[0003] Large conductance calcium activated potassium channels (also BK channels or maxi-K channels) are expressed especially in neurons and smooth muscle cells. Because both of the increase of intracellular calcium concentration and membrane depolarization can activate maxi-K channels, maxi-K channels have been thought to play a pivotal role in regulating voltage-dependent calcium influx. Increase in the intracellular calcium concentration mediates many processes such as release of neurotransmitters, contraction of smooth muscles, cell growth and death, and the like. Actually, the opening of maxi-K channels causes strong membrane hyperpolarization, and inhibits these calcium-induced responses thereby. Accordingly, by inhibiting various depolarization-mediated physiological responses, a substance having an activity of opening maxi-K channels is expected to have potential for the treatment of diseases such as cerebral infarction, subarachnoid hemorrhage, pollakiuria, urinary incontinence, and the like.

[0004] Also, there has been disclosed that a medicine which opens a BK channel can be an agent for treatment of sexual function disorder such as erectile dysfunction, etc. as disclosed in International Publication WO00/34244.

[0005] There have been various reports on a large conductance calcium-activated potassium channel opener. For example, pyrrole derivatives have been disclosed in International Publications WO96/40634, a furan derivative has been disclosed in Japanese Provisional Patent Publication No. 2000-351773, a nitrogen-containing 5-membered derivative in which the nitrogen is substituted by phenyl or benzyl has been disclosed in International Publication WO98/04135 and a diphenyltriazole derivative has been disclosed in Journal of Medicinal Chemistry, vol. 45, pp. 2942-2952, 2002.

[0006] Also, cyclooxygenase-2 inhibitors such as celecoxib, valdecoxib, etc. have been used as a medicine for inflammatory diseases such as rheumatoid arthritis as disclosed in Japanese PCT provisional publications No. 9-506350 and No. 9-500372. However, there is no report of these compounds for use as a large conductance calcium-activated K channel opener.


SUMMARY OF THE INVENTION

[0007] An object of the present invention is to provide an excellent large conductance calcium-activated K channel opener, which is useful for treatment of disorders or diseases such as pollakiuria, urinary incontinence, cerebral infarction, subarachnoid hemorrhage, and the like.




[0008] The present inventors have studied intensively to solve the problems, and as a result, they have found that a certain kind of compounds which have been known as a cyclooxygenase-2 inhibitor have an excellent large conductance calcium-activated K channel opening activity, whereby they have accomplished the present invention.

[0009] That is, the present invention is as mentioned below:

(1) A large conductance calcium-activated K channel opener comprising a compound of the formula (I):




wherein R¹ is a halogen, aminosulfonyl, an alkylsulfonyl or an alkanoylamino sulfonyl; R² is hydrogen or a halogen; R³ and R⁴ may be the same or different from each other and each is hydrogen, a halogen, an alkyl or an alkoxy; Ring A is benzene, pyridine or a cycloalkane, and Ring Q is




where R⁵ is a halogen, an alkyl or a haloalkyl; R⁶ is hydrogen or an alkyl; or R⁵ and R⁶ may be combined to each other to form oxo; R⁷ and R⁸ are hydrogen or may be combined to each other to form oxo; and R⁹ is a carboxyalkyl,

or Ring Q and Ring A may be combined to each other to form a fused ring of the formula:





where X is sulfur or oxygen, and R³, R⁴ and R⁵ have the same meanings as defined above, or a pharmaceutically acceptable salt thereof as an active ingredient.

(2) The large conductance calcium-activated K channel opener as mentioned in the above (1), wherein the opener contains a compound of the formula (II):



wherein R^{1a} is amino, an alkyl or an alkanoylamino; R² is hydrogen or a halogen; R³ and R⁴ may be the same or different from each other and each is hydrogen, a halogen, an alkyl or an alkoxy; Ring A' is benzene or a cycloalkane, and Ring Q' is



where R⁵ is a halogen, an alkyl or a haloalkyl; R⁶ is hydrogen or an alkyl; or R⁵ and R⁶ may be combined to each other to form oxo, or a pharmaceutically acceptable salt thereof as an active ingredient.

(3) The large conductance calcium-activated K channel opener as mentioned in the above (1), wherein the opener contains at least one compound selected from the group consisting of the compounds (1) to (46) mentioned below, or a pharmaceutically acceptable salt thereof as an active ingredient.

(4) The large conductance calcium-activated K channel opener as mentioned in the above (1), wherein the opener contains at least one compound selected from the group consisting of the compounds (1) to (3), (10), (21) to (23), (36), (37) and (43) to (45) mentioned below, or a pharmaceutically acceptable salt thereof as an active ingredient.

(5) A large conductance calcium-activated K channel opener according to any one of the above-mentioned (1) to (4), wherein the opener is an agent for the prophylaxis or treatment of pollakiuria or urinary incontinence.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

[0010] The active ingredient compounds of the present invention have common structural characteristic feature in which they have 4- to 6-membered ring to which two 6-membered rings are substituted at the adjacent portions thereof.

According to this structural characteristic feature, these compounds have an excellent large conductance calcium-activated K channel opening activity.

[0011] In the following, respective groups represented by the respective symbols in the specification are explained.

[0012] The halogen of R¹, R², R³ and R⁵ means fluorine, chlorine, bromine and iodine, and fluorine and chlorine are preferred.

[0013] The alkylsulfonyl of R¹ is exemplified by an alkylsulfonyl having 1 to 4 carbon atoms, specifically methylsulfonyl, ethylsulfonyl, etc., and preferably methylsulfonyl.

[0014] The alkanoylamino of R¹ is exemplified by an alkanoylamino having 2 to 4 carbon atoms, specifically acetylamino, propionylamino, etc. The alkanoylamino portion may form a salt with an alkali metal (such as sodium).

[0015] The alkyl of R³, R⁴, R⁵, R⁶ and R^{1a} is exemplified by an alkyl having 1 to 4 carbon atoms, specifically methyl, ethyl, etc., and preferably methyl.

[0016] The alkoxy of R³ and R⁴ is exemplified by an alkoxy having 1 to 4 carbon atoms, specifically methoxy, ethoxy, etc., and preferably methoxy.

[0017] The cycloalkane of Ring A and Ring A' is exemplified by a cycloalkane having 3 to 7 carbon atoms, specifically cyclopentane, cyclohexane, etc., and preferably cyclohexane.

[0018] The haloalkyl of R⁵ is exemplified by a haloalkyl having 1 to 4 carbon atoms, specifically chloromethyl, trifluoromethyl, etc.


[0019] The carboxyalkyl of R⁹ is exemplified by a carboxyalkyl having 1 to 4 carbon atoms, specifically carboxymethyl.

[0020] The alkanoylamino of R^{1a} is exemplified by an alkanoylamino having 2 to 4 carbon atoms, specifically acetylamino, propionylamino, etc. Moreover, the alkanoylamino portion may form a salt with an alkali metal (such as sodium).

[0021] Examples of pharmaceutically acceptable salts of the compound as an active ingredient may include a salt with an inorganic base such as sodium salt, potassium salt, etc., and a salt with an inorganic acid such as hydrochloride, sulfate, etc.


[0022] The compounds (1) to (18) which are active ingredients of the present invention have the chemical structure as shown in the following Table 1.

Table 1



Compound No.	R ¹	R ²	R ³	R ⁴	Ring Q	Ring A
(1)	4-SO ₂ NH ₂	H	4-CH ₃	H		Phenyl
(2)	4-SO ₂ CH ₃	H	H	H		Phenyl
(3)	4-SO ₂ NH ₂	H	H	H		Phenyl
(4)	4-SO ₂ N (Na) COEt	H	H	H		Phenyl
(5)	4-SO ₂ NH ₂	3-F	H	H		Cyclohexyl
(6)	4-SO ₂ NH ₂	H	4-OCH ₃	3-F		Phenyl
(7)	4-SO ₂ CH ₃	H	3-F	5-F		Phenyl
(8)	4-SO ₂ CH ₃	H	4-F	H		Phenyl
(9)	4-SO ₂ NH ₂	H	4-Cl	H		Phenyl
(10)	4-SO ₂ NH ₂	H	H	H		Phenyl
(11)	4-SO ₂ NH ₂	H	H	H		Phenyl

Table 1 (contd.)



Compound No.	R ¹	R ²	R ³	R ⁴	Ring Q	Ring A
(12)	4-SO ₂ CH ₃	H	4-F	H		Phenyl
(13)	4-SO ₂ NH ₂	H	4-OCH ₃	H		Phenyl
(14)	4-SO ₂ CH ₃	H	4-CH ₃	H		3-Pyridyl
(15)	4-SO ₂ CH ₃	H	H	H		Phenyl
(16)	4-SO ₂ CH ₃	H	H	H		Phenyl
(17)	4-SO ₂ NH ₂	H	4-OCH ₃	5-F		
(18)	4-Cl	H	H	H		Phenyl

[0023] The compounds (1) to (18) can be prepared according to the preparation methods disclosed in the following publications.


- (1) International Publication WO95/15318 pamphlet (Japanese PCT Provisional Patent Publication No. 9-506350),
- (2) International Publication WO95/00501 pamphlet (Japanese PCT Provisional Patent Publication No. 9-500372),
- (3) International Publication WO97/38986 pamphlet (Japanese PCT Provisional Patent Publication No. 2000-509029),
- (4) International Publication WO96/25405 pamphlet (Japanese PCT Provisional Patent Publication No. 11-503722),
- (5) International Publication WO96/19463 pamphlet (Japanese Provisional Patent Publication No. 9-52882),
- (6) International Publication WO00/23426 pamphlet (Japanese PCT Provisional Patent Publication No. 2002-527508),
- (7) European Patent 863134 A (Japanese Provisional Patent Publication No. 10-251220),
- (8) International Publication WO95/11883 pamphlet (Japanese PCT Provisional Patent Publication No. 9-504288),

EP 1 400 243 A1

- (9) International Publication WO95/15316 pamphlet (Japanese PCT Provisional Patent Publication No. 9-506350),
(10) International Publication WO97/29776 pamphlet (Japanese PCT Provisional Patent Publication No. 2000-504723),
5 (11) International Publication WO99/14205 pamphlet (Japanese PCT Provisional Patent Publication No. 2001-516750),
(12) International Publication WO95/30656 pamphlet,
(13) European Patent 799823 A (Japanese Provisional Patent Publication No. 9-323971),
(14) International Publication WO98/03484 pamphlet (Japanese PCT Provisional Patent Publication No. 11-514008),
10 (15) International Publication WO97/36863 pamphlet (Japanese PCT Provisional Patent Publication No. 2000-510444),
(16) International Publication WO96/09304 pamphlet,
(17) International Publication WO96/41626 pamphlet (Japanese PCT Provisional Patent Publication No. 11-507670),
15 (18) Japanese Provisional Patent Publication No. 3-141261.


[0024] Also, the compounds of the formulae (I) and (II) can be prepared according to the preparation methods disclosed in the above-mentioned publications. For example, the following compounds (19) to (46) shown in Table 2 were prepared.

Table 2



Compound No.	R ¹	R ²	R ³	R ⁴	Ring Q	Ring A	Physical properties
(19)	4-SO ₂ NH ₂	H	4-Cl	H		Phenyl	MS·APCI (m/z) : 402 (M+H) ⁺
(20)	4-SO ₂ NHCOCH ₃	H	4-CH ₃	H		Phenyl	MS·APCI (m/z) : 424 (M+H) ⁺
(21)	4-SO ₂ NH ₂	H	4-CH ₃	H		Phenyl	MS·APCI (m/z) : 362/364 (M+H) ⁺
(22)	4-SO ₂ NH ₂	H	4-CH ₃	H		Phenyl	MS·APCI (m/z) : 328 (M+H) ⁺
(23)	4-SO ₂ NH ₂	H	2-CH ₃	H		Phenyl	MS·APCI (m/z) : 382 (M+H) ⁺
(24)	4-SO ₂ NH ₂	H	3-CH ₃	H		Phenyl	MS·APCI (m/z) : 382 (M+H) ⁺
(25)	4-SO ₂ NH ₂	H	2-Cl	H		Phenyl	MS·APCI (m/z) : 402/404 (M+H) ⁺
(26)	4-SO ₂ NH ₂	H	3-Cl	H		Phenyl	MS·APCI (m/z) : 402/404 (M+H) ⁺
(27)	4-SO ₂ NH ₂	H	4-CH ₃	H		Phenyl	MS·APCI (m/z) : 356 (M+H) ⁺

Table 2 (Contd.)














Compound No.	R^1	R^2	R^3	R^4	Ring Q	Ring A	Physical properties
(28)	4-SO ₂ NH ₂	H	4-CH ₃	H		Phenyl	MS·APCI (m/z): 342 (M+H) ⁺
(29)	4-SO ₂ NH ₂	H	4-CH ₃	H		Phenyl	MS·APCI (m/z): 356 (M+H) ⁺
(30)	4-SO ₂ NH ₂	H	H	H		Phenyl	MS·APCI (m/z): 368 (M+H) ⁺
(31)	4-SO ₂ NH ₂	H	2-OCH ₃	H		Phenyl	MS·APCI (m/z): 398 (M+H) ⁺
(32)	4-SO ₂ NH ₂	H	3-OCH ₃	H		Phenyl	MS·APCI (m/z): 398 (M+H) ⁺
(33)	4-SO ₂ NH ₂	H	4-OCH ₃	H		Phenyl	MS·APCI (m/z): 398 (M+H) ⁺
(34)	4-SO ₂ NH ₂	H	3-F	H		Phenyl	MS·APCI (m/z): 386 (M+H) ⁺
(35)	4-SO ₂ NH ₂	H	4-F	H		Phenyl	MS·APCI (m/z): 386 (M+H) ⁺
(36)	4-SO ₂ NH ₂	H	2-F	H		Phenyl	MS·APCI (m/z): 386 (M+H) ⁺
(37)	4-SO ₂ NH ₂	3-OCH ₃	4-OCH ₃	H		Phenyl	MS·APCI (m/z): 428 (M+H) ⁺

Table 2 (Contd.)



Compound No.	R ¹	R ²	R ³	R ⁴	Ring Q	Ring A	Physical properties
(38)	4-SO ₂ CH ₃	H	4-CH ₃	H		Phenyl	MS·APCI (m/z) : 381 (M+H) ⁺
(39)	4-F	H	4-CH ₃	H		Phenyl	MS·APCI (m/z) : 321 (M+H) ⁺
(40)	3-Cl	H	4-CH ₃	H		Phenyl	MS·APCI (m/z) : 337/339 (M+H) ⁺
(41)	2-Cl	H	4-CH ₃	H		Phenyl	MS·APCI (m/z) : 337/339 (M+H) ⁺
(42)	4-Cl	H	4-CH ₃	H		Phenyl	MS·APCI (m/z) : 337/339 (M+H) ⁺
(43)	4-SO ₂ NH ₂	H	4-CH ₃	3-CH ₃		Phenyl	MS·APCI (m/z) : 396 (M+H) ⁺
(44)	4-SO ₂ NH ₂	H	H	H		3-Pyridyl	MS·APCI (m/z) : 369 (M+H) ⁺
(45)	4-SO ₂ NH ₂	H	4-Br	H		Phenyl	MS·APCI (m/z) : 393/395 (M+H) ⁺
(46)	4-SO ₂ CH ₃	H	H	H		Phenyl	MS·APCI (m/z) : 314 (M+H) ⁺

[0025] The active ingredient compounds of the present invention or pharmaceutically acceptable salts thereof can be administered orally or parenterally with a pharmaceutically acceptable carrier or diluent and used as common phar-

maceutical preparations such as tablets, granules, capsules, powders, injection and inhalants.

[0026] As a pharmaceutically acceptable carrier for a preparation of oral administration, there may be mentioned a material commonly used, for example, a binder (such as syrup, Gum Arabic, gelatin, sorbit, tragacanth and polyvinyl pyrrolidone), an excipient (such as lactose, sugar, corn starch, potassium phosphate, sorbit and glycine), a lubricant (such as magnesium stearate, talc, polyethylene glycol and silica), a disintegrator (such as potato starch) and a humectant (such as anhydrous lauryl sodium sulfate).

[0027] On the other hand, when the active ingredient compounds of the present invention are administered parenterally, it may be formulated into the form of an injection or a drip infusion by using distilled water for injection, physiological saline, an aqueous glucose solution and the like, or a suppository.

[0028] A dose of the active ingredient compounds of the present invention or pharmaceutically acceptable salts thereof may vary depending on an administration method, an age, weight or conditions of a patient, or a kind or degree of disease, and is generally about 0.01 to 50 mg/kg per day, more preferably about 0.1 to 30 mg/kg per day.

[0029] The active ingredient compounds of the present invention or pharmaceutically acceptable salts thereof have an excellent large conductance calcium-activated K channel opening activity and hyperpolarizes a membrane electric potential of cells, so that it may be used for the prophylactic, relief and/or treatment of, for example, hypertension, premature birth, irritable bowel syndrome, chronic heart failure, angina, cardiac infarction, cerebral infarction, subarachnoid hemorrhage, cerebral vasospasm, cerebral hypoxia, peripheral blood vessel disorder, anxiety, male-pattern baldness, erectile dysfunction, diabetes, diabetic peripheral nerve disorder, other diabetic complication, sterility, urolithiasis and pain accompanied thereby, pollakiuria, urinary incontinence, nocturnal enuresis, asthma, chronic obstructive pulmonary disease (COPD), cough accompanied by asthma or chronic obstructive pulmonary disease (COPD), cerebral apoplexy, cerebral ischemia, traumatic encephalopathy, and the like.

BEST MODE OF EMBODIMENT OF THE INVENTION

[0030] In the following, the large conductance calcium-activated K channel opener of the present invention is explained in more detail by referring to Experimental examples, but the present invention is not limited by these.

Experimental example 1

Relaxation effect on potassium-induced contraction of isolated rabbit urinary bladder

[0031] Urinary bladder was isolated from rabbits (weight: 2.0 to 3.5 kg) and immediately immersed in ice-cold Krebs-bicarbonate solution (in mM: 118 NaCl, 4.7 KCl, 2.55 CaCl₂, 1.18 MgSO₄, 1.18 KH₂PO₄, 24.88 NaHCO₃, and 11.1 glucose) to remove peripheral binding tissues. The urinary bladder was cut into longitudinal strips (about 5 mm length, 3 to 4 mm width) after mucosal layer was removed.

[0032] Preparations were mounted in organ baths containing Krebs solution maintained at 37°C and gassed with 95% O₂/5% CO₂.

[0033] Accordingly, preparations were stretched with an initial tension of 2.0 g, and changes in isometric tension were measured by force-displacement transducer. The preparations were pre-contracted by changing organ-bath solution into high-K⁺ (30 mM) Krebs solution.

[0034] After stable tension was obtained, compounds were added into organ baths cumulatively (10⁻⁸ M-10⁻⁴ M). The effects of compounds were expressed as a percentage of the maximum relaxation produced by 0.1 mM papaverine. 50% relaxation concentration (IC₅₀) was calculated and IC₅₀ value (μM) of the compounds of the present invention was shown in the following Table 3.

Table 3

Compound to be tested	IC ₅₀ (μM)
Compound (1)	9.24
Compound (3)	1.53
Compound (23)	0.11

Experimental example 2

Inhibitory effect on the rhythmic bladder contractions induced by substance P in anesthetized rats

[0035] For the experiments, Sprague-Dawley female rats (9 to 12 weeks old) weighing between 200 to 300 g were

used. After urethane anesthetization (subcutaneously administered with a dose of 1.2 g/kg), cannulae were placed in both right and left femoral veins. One intravenous catheter was used for administration of compounds, and the other was for the substance P (0.33 $\mu\text{g/kg/min}$) infusion. We also cannulated into ureter to pass urine. Polyethylene catheters were inserted into carotid artery for continuous monitoring of arterial blood pressure and heart rate. For continuous infusion, transurethral bladder catheter was inserted into the bladder through the urethra and tied in place by a ligature around the urethral orifice. One end of the catheter was attached to a pressure transducer in order to measure intravesical pressure. The other end of the catheter was used for infusion of saline into the bladder. After stabilization of blood pressure and heart rate and after the bladder was emptied, cystometry was performed by filling the bladder slowly with about 0.6 ml of saline. After about 10 minutes, intravenous infusion of substance P (0.33 $\mu\text{g/kg/min}$) was started for stabilization of the micturition reflex. Compounds were administered after stable rhythmic bladder contraction was obtained over 15 minutes. All compounds were dissolved or suspended in saline containing 0.5% Tween 80 for intravenous administration (0.1 ml/kg). The rhythmic contraction frequency and the intravesical pressure were observed for 35 minutes after administration of the test compound.

[0036] As a result, the compounds of the present invention decreased the frequency of bladder rhythmic contraction without changing the amplitude of contraction. Also, we determined a time (minute) during which the frequency of the rhythmic contraction had been completely inhibited by administering 0.25 mg/kg of the compound. A 100% inhibition time (minute) of the selected compounds of the present invention is shown in the following Table 4.


Table 4

Compound to be tested	Time (min)
Compound (1)	11
Compound (3)	16.7
Compound (23)	8

[0037] Also, it was shown that an inhibitory activity of causing the bladder rhythmic contraction had been weakened by pre-administration of iberiotoxin (0.15 mg/kg, intravenous administration) which is a specific inhibitor for a large conductance calcium-activated K channel opener. Accordingly, it has been shown that the active ingredient of the present invention is effective for a prophylaxis or treatment agent for pollakiuria or urinary incontinence through a large conductance calcium-activated K channel opening activity.


[0038] Of the compounds shown by the formulae (I) and (II), other preferred compounds may be mentioned in the following Table 5.

Table 5




Compound No.	R^1	R^2	R^3	R^4	Ring Q	Ring A
(47)	4-SO ₂ CH ₃	H	4-F	H		Phenyl
(48)	4-SO ₂ CH ₃	H	4-F	3-F		Phenyl
(49)	4-SO ₂ CH ₃	H	4-Cl	H		Phenyl
(50)	4-SO ₂ CH ₃	H	4-OCH ₃	H		Phenyl
(51)	4-SO ₂ CH ₃	H	2-F	H		Phenyl

Table 5 (Contd.)




Compound No.	R ¹	R ²	R ³	R ⁴	Ring Q	Ring A
(52)	4-SO ₂ NH ₂	H	4-Cl	3-Cl		Phenyl
(53)	4-SO ₂ NH ₂	H	4-F	3-F		Phenyl
(54)	4-SO ₂ NH ₂	H	4-OCH ₃	3-Cl		Phenyl
(55)	4-SO ₂ NH ₂	H	4-F	H		Phenyl
(56)	4-SO ₂ NH ₂	H	4-Cl	H		Phenyl
(57)	4-SO ₂ NH-Ac	H	H	H		Phenyl
(58)	4-SO ₂ CH ₃	H	H	H		Cyclohexyl
(59)	4-SO ₂ NH ₂	H	H	H		Cyclohexyl
(60)	4-SO ₂ NH-CH ₃	H	H	H		Cyclohexyl
(61)	4-SO ₂ NH ₂	3-F	H	H		Cyclohexyl
(62)	4-SO ₂ CH ₃	3-F	H	H		Cyclohexyl
(63)	4-SO ₂ CH ₃	H	4-F	H		Phenyl
(64)	4-SO ₂ NH ₂	H	H	H		Phenyl
(65)	4-SO ₂ NH ₂	H	H	H		Phenyl

Table 5 (Contd.)




Compound No.	R^1	R^2	R^3	R^4	Ring Q	Ring A
(66)	4-SO ₂ NH ₂	H	H	H		Phenyl
(67)	4-SO ₂ NH ₂	3-CH ₃	H	H		Phenyl
(68)	4-SO ₂ NH ₂	H	H	H		Phenyl
(69)	4-SO ₂ CH ₃	H	4-CH ₃	H		Phenyl
(70)	4-SO ₂ NH ₂	H	3-F	H		Phenyl
(71)	4-SO ₂ CH ₃	H	4-F	H		Phenyl
(72)	4-SO ₂ CH ₃	H	4-F	H		Phenyl
(73)	4-SO ₂ NH ₂	H	4-F	H		Phenyl
(74)	4-SO ₂ NH ₂	H	4-CH ₃	H		Phenyl
(75)	4-SO ₂ NH ₂	H	4-Cl	H		3-Pyridyl
(76)	4-SO ₂ CH ₃	H	4-Et	H		Phenyl
(77)	4-SO ₂ NH ₂	H	3-CH ₃	H		Phenyl
(78)	4-SO ₂ NH ₂	H	4-CH ₃	H		Phenyl

Table 5 (Contd.)




Compound No.	R^1	R^2	R^3	R^4	Ring Q	Ring A
(79)	4-SO ₂ NH ₂	H	4-OCH ₃	H		Phenyl
(80)	4-SO ₂ NH ₂	H	4-CH ₃	3-F		Phenyl
(81)	4-SO ₂ NH ₂	H	4-F	3-F		Phenyl
(82)	4-SO ₂ CH ₃	H	H	H		Phenyl
(83)	4-SO ₂ CH ₃	H	4-F	H		Phenyl
(84)	4-SO ₂ CH ₃	H	H	H		3-Pyridyl
(85)	4-SO ₂ CH ₃	H	4-F	H		3-Pyridyl
(86)	4-SO ₂ CH ₃	H	3-Cl	H		3-Pyridyl
(87)	4-SO ₂ CH ₃	H	4-OCH ₃	H		Phenyl
(88)	4-SO ₂ CH ₃	H	H	H		Phenyl
(89)	4-SO ₂ CH ₃	H	4-Cl	H		Phenyl
(90)	4-SO ₂ CH ₃	H	H	H		Phenyl
(91)	4-SO ₂ CH ₃	H	H	H		2-Pyridyl
(92)	4-SO ₂ CH ₃	H	4-OCH ₃	H		3-Pyridyl

Table 5 (Contd.)




Compound No.	R ¹	R ²	R ³	R ⁴	Ring Q	Ring A
(93)	4-SO ₂ CH ₃	H	H	H		Phenyl
(94)	4-SO ₂ NH ₂	H	H	H		Phenyl
(95)	4-SO ₂ NH ₂	H	4-F	H		Phenyl
(96)	4-SO ₂ NH ₂	H	4-F	H		3-Pyridyl
(97)	4-SO ₂ NH ₂	H	4-Cl	H		Phenyl
(98)	4-SO ₂ NH ₂	H	H	H		3-Pyridyl
(99)	4-SO ₂ NH ₂	H	4-CH ₃	6-CH ₃		3-Pyridyl
(100)	4-SO ₂ CH ₃	H	4-F	H		Phenyl
(101)	4-SO ₂ NH ₂	H	4-F	H		Phenyl
(102)	4-SO ₂ CH ₃	H	4-Cl	H		Phenyl
(103)	4-SO ₂ NH ₂	H	4-Cl	H		Phenyl
(104)	4-SO ₂ CH ₃	H	4-F	H		Phenyl
(105)	4-SO ₂ NH ₂	H	4-F	H		Phenyl
(106)	4-SO ₂ CH ₃	H	4-Cl	H		Phenyl
(107)	4-SO ₂ NH ₂	H	4-Cl	H		Phenyl

Table 5 (Contd.)



Compound No.	R ¹	R ²	R ³	R ⁴	Ring Q	Ring A
(108)	4-SO ₂ NH ₂	H	4-F	H		Phenyl
(109)	4-SO ₂ NH ₂	H	4-F	H		Phenyl
(110)	4-SO ₂ CH ₃	H	H	H		Phenyl
(111)	4-SO ₂ CH ₃	H	4-F	H		Phenyl
(112)	4-SO ₂ CH ₃	H	4-OCH ₃	H		Phenyl
(113)	4-SO ₂ CH ₃	H	4-CH ₃	H		Phenyl
(114)	4-SO ₂ NH ₂	H	H	H		Phenyl
(115)	4-SO ₂ NH ₂	H	4-F	H		Phenyl
(116)	4-SO ₂ NH ₂	H	4-OCH ₃	H		Phenyl
(117)	4-SO ₂ NH ₂	H	4-CH ₃	H		Phenyl
(118)	4-SO ₂ NH ₂	H	4-F	3-F		Phenyl
(119)	4-SO ₂ NH ₂	H	4-Cl	H		Phenyl
(120)	4-SO ₂ NH ₂	H	4-CH ₃	3-CH ₃		Phenyl
(121)	4-SO ₂ NH ₂	H	4-F	H		Phenyl
(122)	4-SO ₂ NH ₂	H	4-Cl	H		Phenyl

Table 5 (Contd.)



Compound No.	R ¹	R ²	R ³	R ⁴	Ring Q	Ring A
(123)	4-SO ₂ NH ₂	H	H	H		Phenyl
(124)	4-SO ₂ CH ₃	H	4-F	H		Phenyl
(125)	4-SO ₂ CH ₃	H	4-Cl	H		Phenyl

INDUSTRIAL APPLICABILITY

[0039] The active ingredient compounds of the present invention or pharmaceutically acceptable salts thereof have an excellent large conductance calcium-activated K channel opening activity, so that it is used for the prophylactic, relief and/or treatment of hypertension, premature birth, irritable bowel syndrome, chronic heart failure, angina, cardiac infarction, cerebral infarction, subarachnoid hemorrhage, cerebral vasospasm, cerebral hypoxia, peripheral blood vessel disorder, anxiety, male-pattern baldness, erectile dysfunction, diabetes, diabetic peripheral nerve disorder, other diabetic complication, sterility, urolithiasis and pain accompanied thereby, pollakiuria, urinary incontinence, nocturnal enuresis, asthma, chronic obstructive pulmonary disease (COPD), cough accompanied by asthma or chronic obstructive pulmonary disease (COPD), cerebral apoplexy, cerebral ischemia, traumatic encephalopathy, and the like. Also, the active ingredient compounds of the present invention or pharmaceutically acceptable salts thereof have a low toxicity, so that it has high safety as a medicine.

Claims

1. Use of a compound of formula (I):



15




45

55





or a pharmaceutically acceptable salt thereof, for the manufacture of a medicament for the prophylaxis or treatment of a disorder or disease responsive to opening of the large conductance calcium-activated K channel.

2. Use according to claim 1, wherein the compound of formula (I) is a compound of formula (II):



wherein R^{1a} is amino, an alkyl or an alkanoylamino; R^2 is hydrogen or a halogen; R^3 and R^4 may be the same or different from each other and each is hydrogen, a halogen, an alkyl or an alkoxy; Ring A' is benzene or a cycloalkane, and Ring Q' is



where R^5 is a halogen, an alkyl or a haloalkyl; R^6 is hydrogen or an alkyl; or R^5 and R^6 may be combined to each other to form oxo.

3. Use according to claim 1, wherein the compound of formula (I) is a compound selected from the group consisting of:

- (1) celecoxib,
- (2) rofecoxib,
- (3) valdecoxib,
- (4) parecoxib,
- (5) tilmacoxib,
- (6) 4-(4-chloro-5-(3-fluoro-4-methoxyphenyl)imidazol-1-yl)benzenesulfonamide,
- (7) 2-(3,5-difluorophenyl)-3-(4-methylsulfonylphenyl)-2-cyclopenten-1-one,
- (8) 1-fluoro-4-(2-(4-methylsulfonylphenyl)-1-cyclopenten-1-yl)benzene,
- (9) 4-(5-(4-chlorophenyl)-3-trifluoromethyl-1H-pyrazol-1-yl)benzenesulfonamide,
- (10) 4-(2-methyl-4-phenyloxazol-5-yl)benzenesulfonamide,
- (11) 4-(2-oxo-3-phenyl-2,3-dihydroxazol-4-yl)benzene-sulfonamide,
- (12) 1-(3,3-dimethyl-5-(4-methylsulfonylphenyl)cyclopenta-1,4-dien-1-yl)-4-fluorobenzene,

- (13) 4-(2-(4-methoxyphenyl)-4-methylpyrrol-1-yl)benzene-sulfonamide,
 (14) etoricoxib,
 (15) 4,4-dimethyl-2-phenyl-3-(4-methylsulfonylphenyl)cyclo-butanone,
 (16) 5-(4-methylsulfonylphenyl)-6-phenyl[1,3]thiazole[3,2-b][1,2,4]triazole,
 5 (17) 4-(6-fluoro-7-methoxy-3-trifluoromethylisothio-chromeno[4,3-c]pyrazol-1(5H)-yl)benzenesulfonamide,
 (18) licofelone,
 (19) 4-[5-(4-chlorophenyl)-3-trifluoromethyl-1H-pyrazol-1-yl]benzenesulfonamide,
 (20) N-acetyl-4-[5-(4-methylphenyl)-3-trifluoromethyl-1H-pyrazol-1-yl]benzenesulfonamide,
 (21) 4-[5-(4-methylphenyl)-3-chloromethyl-1H-pyrazol-1-yl]-benzenesulfonamide,
 10 (22) 4-[5-(4-methylphenyl)-3-methyl-1H-pyrazol-1-yl]benzenesulfonamide,
 (23) 4-[5-(2-methylphenyl)-3-trifluoromethyl-1H-pyrazol-1-yl]benzenesulfonamide,
 (24) 4-[5-(3-methylphenyl)-3-trifluoromethyl-1H-pyrazol-1-yl]benzenesulfonamide,
 (25) 4-[5-(2-chlorophenyl)-3-trifluoromethyl-1H-pyrazol-1-yl]benzenesulfonamide,
 (26) 4-[5-(3-chlorophenyl)-3-trifluoromethyl-1H-pyrazol-1-yl]benzenesulfonamide,
 15 (27) 4-[5-(4-methylphenyl)-3-n-propyl-1H-pyrazol-1-yl]benzenesulfonamide,
 (28) 4-[5-(4-methylphenyl)-3-ethyl-1H-pyrazol-1-yl]benzene-sulfonamide,
 (29) 4-[5-(4-methylphenyl)-3-isopropyl-1H-pyrazol-1-yl]benzenesulfonamide,
 (30) 4-[5-phenyl-3-trifluoromethyl-1H-pyrazol-1-yl]benzene-sulfonamide,
 (31) 4-[5-(2-methoxyphenyl)-3-trifluoromethyl-1H-pyrazol-1-yl]benzenesulfonamide,
 20 (32) 4-[5-(3-methoxyphenyl)-3-trifluoromethyl-1H-pyrazol-1-yl]benzenesulfonamide,
 (33) 4-[5-(4-methoxyphenyl)-3-trifluoromethyl-1H-pyrazol-1-yl]benzenesulfonamide,
 (34) 4-[5-(3-fluorophenyl)-3-trifluoromethyl-1H-pyrazol-1-yl]benzenesulfonamide,
 (35) 4-[5-(4-fluorophenyl)-3-trifluoromethyl-1H-pyrazol-1-yl]benzenesulfonamide,
 (36) 4-[5-(2-fluorophenyl)-3-trifluoromethyl-1H-pyrazol-1-yl]benzenesulfonamide,
 25 (37) 4-[5-(3,4-dimethoxyphenyl)-3-trifluoromethyl-1H-pyrazol-1-yl]benzenesulfonamide,
 (38) 5-(4-methylphenyl)-1-(4-methylsulfonylphenyl)-3-trifluoromethyl-1H-pyrazole,
 (39) 5-(4-methylphenyl)-1-(4-fluorophenyl)-3-trifluoromethyl-1H-pyrazole,
 (40) 5-(4-methylphenyl)-1-(3-chlorophenyl)-3-trifluoromethyl-1H-pyrazole,
 (41) 5-(4-methylphenyl)-1-(2-chlorophenyl)-3-trifluoromethyl-1H-pyrazole,
 30 (42) 5-(4-methylphenyl)-1-(4-chlorophenyl)-3-trifluoromethyl-1H-pyrazole,
 (43) 4-[5-(3,4-dimethylphenyl)-3-trifluoromethyl-1H-pyrazol-1-yl]benzenesulfonamide,
 (44) 4-[5-(3-pyridyl)-3-trifluoromethyl-1H-pyrazol-1-yl]-benzenesulfonamide,
 (45) 4-[5-methyl-3-(4-bromophenyl)isoxazol-4-yl]benzenesulfonamide, and
 (46) 5-methyl-3-phenyl-4-(4-methylsulfonylphenyl)isoxazole.

4. Use according to claim 1, wherein the compound of formula (I) is a compound selected from the group consisting of:

- (1) celecoxib,
 (2) rofecoxib,
 40 (3) valdecoxib,
 (10) 4-(2-methyl-4-phenyloxazol-5-yl)benzenesulfonamide,
 (21) 4-[5-(4-methylphenyl)-3-chloromethyl-1H-pyrazol-1-yl]benzenesulfonamide,
 (22) 4-[5-(4-methylphenyl)-3-methyl-1H-pyrazol-1-yl]benzenesulfonamide,
 (23) 4-[5-(2-methylphenyl)-3-trifluoromethyl-1H-pyrazol-1-yl]benzenesulfonamide,
 45 (36) 4-[5-(2-fluorophenyl)-3-trifluoromethyl-1H-pyrazol-1-yl]benzenesulfonamide,
 (37) 4-[5-(3,4-dimethoxyphenyl)-3-trifluoromethyl-1H-pyrazol-1-yl]benzenesulfonamide,
 (43) 4-[5-(3,4-dimethylphenyl)-3-trifluoromethyl-1H-pyrazol-1-yl]benzenesulfonamide,
 (44) 4-[5-(3-pyridyl)-3-trifluoromethyl-1H-pyrazol-1-yl]-benzenesulfonamide, and
 50 (45) 4-[5-methyl-3-(4-bromophenyl)isoxazol-4-yl]benzene-sulfonamide.

5. Use according to any one of claims 1 to 4, wherein the disorder or disease responsive to opening of the large conductance calcium-activated K channel is hypertension, premature birth, irritable bowel syndrome, chronic heart failure, angina, cardiac infarction, cerebral infarction, subarachnoid hemorrhage, cerebral vasospasm, cerebral hypoxia, peripheral blood vessel disorder, anxiety, male-pattern baldness, erectile dysfunction, diabetes, diabetic peripheral nerve disorder, other diabetic complication, sterility, urolithiasis and pain accompanied thereby, pollakiuria, urinary incontinence, nocturnal enuresis, asthma, chronic obstructive pulmonary disease (COPD), cough accompanied by asthma or chronic obstructive pulmonary disease (COPD), cerebral apoplexy, cerebral ischemia or traumatic encephalopathy.

EP 1 400 243 A1

6. Use according to any one of claims 1 to 4, wherein the disorder or disease responsive to opening of the large conductance calcium-activated K channel is pollakiuria or urinary incontinence.

5

10

15

20

25

30

35

40

45

50

55



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 03 25 5860

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
X	US 6 440 963 B1 (TESTA RODOLFO ET AL) 27 August 2002 (2002-08-27) * claims 4,14,71 * * column 16 * * column 14, line 38-50 * ---	1-6	A61K31/415 A61K31/33 A61P29/00
X	WO 98 47509 A (CONNOLLY DANIEL T ;SEARLE & CO (US); SEIBERT KAREN (US); RONIKER B) 29 October 1998 (1998-10-29) * claims 1,2,5,8 * ---	1-5	
X	WO 02 40052 A (HEDENSTIERNA GOERAN ;CHEN LUNI (SE); AGA AB (SE)) 23 May 2002 (2002-05-23) * claims 1-3,8 * ---	1-5	
X	EP 1 064 964 A (PFIZER PROD INC) 3 January 2001 (2001-01-03) * page 5, line 54 - page 6, line 33 * * claims 1,2,5-8 * ---	1-5	
X	WO 01 28548 A (WILLERSON JAMES T ;UNIV TEXAS (US); DELGADO REYNOLDS M III (US); T) 26 April 2001 (2001-04-26) * claims 1-3 * -----	1-5	<div>TECHNICAL FIELDS SEARCHED (Int.Cl.7)</div> <div>A61K</div>
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 10 December 2003	Examiner Collura, A
<div>CATEGORY OF CITED DOCUMENTS</div> <div> X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document </div>			

EPO FORM 1500 (03.02) (104001)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 03 25 5860

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

10-12-2003

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 6440963	B1	27-08-2002	IT M120010733 A1	07-10-2002
			WO 02080927 A1	17-10-2002
WO 9847509	A	29-10-1998	AU 745797 B2	28-03-2002
			AU 7466298 A	13-11-1998
			BG 103803 A	31-05-2000
			BR 9808932 A	01-08-2000
			CN 1253502 T	17-05-2000
			EE 9900517 A	15-06-2000
			EE 200300169 A	16-06-2003
			EP 0979077 A1	16-02-2000
			HU 0001777 A2	28-05-2001
			JP 2001527542 T	25-12-2001
			NO 995077 A	17-12-1999
			NZ 500141 A	31-05-2002
			PL 337098 A1	31-07-2000
			SK 138799 A3	18-01-2001
			TR 9902545 T2	21-01-2000
			WO 9847509 A1	29-10-1998
			US 2002035156 A1	21-03-2002
			ZA 9803249 A	19-04-1999
WO 0240052	A	23-05-2002	AU 1451902 A	27-05-2002
			CA 2429224 A1	23-05-2002
			EE 200300226 A	15-08-2003
			EP 1339428 A1	03-09-2003
			WO 0240052 A1	23-05-2002
EP 1064964	A	03-01-2001	BR 0002953 A	30-01-2001
			CA 2313131 A1	30-12-2000
			EP 1064964 A2	03-01-2001
			JP 2001039872 A	13-02-2001
			US 2002061918 A1	23-05-2002
			US 6413965 B1	02-07-2002
WO 0128548	A	26-04-2001	AU 8026200 A	30-04-2001
			WO 0128548 A1	26-04-2001
			US 6323226 B1	27-11-2001

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82